


Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

## Neurobiology of Aging

journal homepage: [www.elsevier.com/locate/neuaging.org](http://www.elsevier.com/locate/neuaging.org)

# Previous estradiol treatment during midlife maintains transcriptional regulation of memory-related proteins by ER $\alpha$ in the hippocampus in a rat model of menopause

Nina E. Baumgartner<sup>a,b,\*</sup>, Katelyn L. Black<sup>a,b</sup>, Shannon M. McQuillen<sup>a,b</sup>, Jill M. Daniel<sup>a,b,c</sup>

<sup>a</sup> Brain Institute, Tulane University, New Orleans, LA

<sup>b</sup> Neuroscience Program, Tulane University, New Orleans, LA

<sup>c</sup> Psychology Department, Tulane University, New Orleans, LA

## ARTICLE INFO

## Article history:

Received 23 February 2021

Revised 27 May 2021

Accepted 30 May 2021

Available online 5 June 2021

## Keywords:

Estrogen

Estrogen receptor

Hippocampus

Menopause

BDNF

ChAT

## ABSTRACT

Previous midlife estradiol treatment, like continuous treatment, improves memory and results in lasting increases in hippocampal levels of estrogen receptor (ER)  $\alpha$  and ER-dependent transcription in ovariectomized rodents. We hypothesized that previous and continuous midlife estradiol act to specifically increase levels of nuclear ER $\alpha$ , resulting in transcriptional regulation of proteins that mediate estrogen effects on memory. Ovariectomized middle-aged rats received estradiol or vehicle capsule implants. After 40 days, rats initially receiving vehicle received another vehicle capsule (ovariectomized controls). Rats initially receiving estradiol received either another estradiol (continuous estradiol) or a vehicle (previous estradiol) capsule. One month later, hippocampi were dissected and processed. Continuous and previous estradiol increased levels of nuclear, but not membrane or cytosolic ER $\alpha$  and had no effect on *Esr1*. Continuous and previous estradiol impacted gene expression and/or protein levels of mediators of estrogenic action on memory including ChAT, BDNF, and PSD-95. Findings demonstrate a long-lasting role for hippocampal ER $\alpha$  as a transcriptional regulator of memory following termination of previous estradiol treatment in a rat model of menopause.

© 2021 The Author(s). Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## 1. Introduction

Decades of research support the idea that estrogens play an important role in modulating memory in the aging female brain (Koebele and Bimonte-Nelson, 2017; Luine and Frankfurt, 2020). Declines in ovarian hormones following menopause coincide with increased risk of pathological and nonpathological cognitive decline. Estrogens administered near the onset of loss of ovarian function have been shown to improve cognition in humans and rodents and enhance function of the hippocampus, a brain region crucial for memory (Daniel et al., 2015; Maki et al., 2011). However, due to serious health risks of long-term estrogen use (Chen et al., 2006), current guidelines recommend that individu-

als who do choose to use estrogens to treat menopause symptoms do so for only a few years near menopause (Santen et al., 2010).

In an aging ovariectomized rodent model of menopause, our lab has shown that short-term estrogen use during midlife has lasting effects on the brain and cognition. Forty days of estradiol exposure immediately following midlife ovariectomy resulted in enhanced performance on the hippocampal-dependent radial arm maze up to 7 months after estradiol treatment had been ended, effects that were comparable to ongoing estradiol treatment (Rodgers et al., 2010). This initial finding in our lab demonstrated that estrogens administered for a short-term period immediately after the loss of ovarian function can have lasting benefits for memory similar to those exerted by ongoing estradiol exposure. Since then, we have replicated this lasting impact of previous exposure to midlife estradiol on memory in rats (Black et al., 2018; Black et al., 2016; Witty et al., 2013) and in mice (Pollard et al., 2018). Evidence in nonhuman primates receiving short-term estrogen use in midlife shows similar results. Ovariectomized rhesus monkeys that re-

\* Corresponding author at: Tulane University, 201 Flower Hall, New Orleans, LA 70118, Phone: (504) 862-3301.

E-mail address: [nbaumgar@tulane.edu](mailto:nbaumgar@tulane.edu) (N.E. Baumgartner).

ceived 11 months of cyclic estradiol injections displayed enhanced performance on a memory task one year after termination of hormone treatment (Baxter et al., 2018). Finally, in women who undergo surgical menopause (oophorectomy) earlier in life than natural menopause, estrogen treatment until the time at which natural menopause would occur is recommended and is associated with decreased risk of cognitive impairment later in life (Bove et al., 2014; Rocca et al., 2014). Collectively, these studies demonstrate across multiple species that exposure to estrogens immediately following loss of ovarian function can improve cognition long after hormone treatment has ended.

The ability of previous midlife estradiol exposure to enhance cognitive aging long-term is related to its ability to impact levels of estrogen receptors in the hippocampus. Estrogens exert their effects on the brain by acting on estrogen receptors, including the classic nuclear steroid receptor estrogen receptors (ER)  $\alpha$ . Previous midlife estradiol exposure resulted in lasting increases in hippocampal levels of ER $\alpha$  eight months after termination of estradiol treatment (Rodgers et al., 2010). Subsequent work in our lab and others suggest a causal relationship between increased hippocampal ER $\alpha$  and memory. For instance, increasing hippocampal ER $\alpha$  using lenti-viral vectors enhances performance on the radial arm maze in aged ovariectomized rats (Witty et al., 2012). Additionally, pharmacologically antagonizing ER $\alpha$  using ICI 182780 prevents the memory benefits shown previously with our short-term estradiol model in aged ovariectomized rats (Black et al., 2016). Finally, certain polymorphisms in *Esr1*, the gene that encodes for ER $\alpha$ , may impact cognitive function (Ma et al., 2014; Yaffe et al., 2009) and increase risk of Alzheimer's disease in postmenopausal women (Ma et al., 2009; Ryan et al., 2014), demonstrating that ER $\alpha$  can impact memory long after cessation of ovarian function.

Sustained increases in hippocampal ER $\alpha$  levels following previous midlife estradiol exposure can have long-lasting impacts on hippocampal function. Activation of ER $\alpha$  can lead to a wide range of changes within a cell, but its actions are often classified into two categories: genomic and nongenomic. The genomic actions of ER $\alpha$  are the classic steroid hormone receptor actions that involve the receptor acting as a transcription factor at estrogen response elements (EREs) to promote genomic changes within a cell (Klinge, 2001). Consistent with the observed impacts on hippocampal memory, we have also shown that previous exposure to estradiol following ovariectomy results in lasting increases in ERE-dependent transcriptional activity in the hippocampus of ovariectomized ERE-luciferase reporter mice (Pollard et al., 2018). Currently, it remains unknown which specific ERE-dependent genes are impacted by previous midlife estradiol exposure that could be involved with the effects of this hormone treatment paradigm on memory. Three potential genes that contain an ERE sequence and are known to regulate memory include *Esr1* (Castles et al., 1997), *Bdnf* (Sohrabji et al., 1995), and *Chat* (Hyder et al., 1999). Whereas *Esr1* is the gene that transcribes ER $\alpha$ , the proteins transcribed by both *Bdnf* and *Chat* are closely associated with the effects of ER $\alpha$  on hippocampal function (Kőszegi et al., 2011; Scharfman and MacLuskey, 2005). Brain-derived neurotrophic factor (BDNF) is involved in hippocampal neurogenesis and neuroprotection in the aging brain (Pencea et al., 2001; Sohrabji and Lewis, 2006). Choline acetyltransferase (ChAT) is the synthesizing enzyme for acetylcholine, a neurotransmitter closely associated with the actions of estrogen in the hippocampus (Gibbs, 1997; Luine, 1985).

In addition to traditional genomic actions associated with nuclear steroid receptors, membrane localized ER $\alpha$  is also able to impact memory in the hippocampus by acting through rapid nongenomic mechanisms. Membrane ER $\alpha$  has been shown to rapidly activate multiple intracellular signaling pathways that influence hippocampal dependent cognition [for review, see (Foster, 2012)].

Rapid effects of estradiol administration on cellular signaling pathways has been implicated in synaptic transmission (Fugger et al., 2001), cell excitability (Kumar and Foster, 2002), NMDA receptor function (Bi et al., 2003), long-term potentiation (Foy et al., 2008), and rapid changes in expression of synaptic proteins including postsynaptic density protein 95 (PSD-95), a protein crucial for stabilizing synaptic changes during long-term potentiation (Akama and McEwen, 2003; Murakami et al., 2015). Because these different actions of ER $\alpha$  are associated with specific locations of the receptor, the subcellular distribution of ER $\alpha$  dictates its function. Currently, it is unknown where the observed increase in ER $\alpha$  following previous midlife estradiol exposure occurs within hippocampal cells.

The overall goal of the current work was to determine mechanisms by which previous exposure to estradiol in midlife, acting through its ability to increase levels of ER $\alpha$ , is able to maintain hippocampal function. To do so we compared impacts of previous estradiol treatment to ongoing estradiol and ovariectomized control treatments on hippocampal gene and protein expression of estrogen-sensitive genes and on the subcellular distribution of hippocampal ER $\alpha$  protein levels. Specifically, we measured gene expression and corresponding protein levels of three estrogen-sensitive genes that contain ERE sequences (*Esr1/ER $\alpha$* , *Bdnf/BDNF*, *Chat/ChAT*) and one estrogen-sensitive gene without a known ERE sequence but associated with the actions of membrane-bound ER $\alpha$  (*Dlg4/PSD-95*). Subcellular fractionation was performed before measuring ER $\alpha$  protein levels in order to determine the subcellular localization of the receptor in hippocampal cells following previous exposure to estradiol in midlife.

## 2. Materials and methods

### 2.1. Subjects

Middle-aged female Long-Evans hooded rats, retired breeders (~11 months of age), were purchased from Envigo. Long-Evans rats at this age are considered middle-aged, a period characterized by irregular or prolonged estrous cycles (Chakraborty and Gore 2004), which is consistent with our own observations (Daniel et al., 2006). Animal care was in accordance with guidelines set by the National Institute of Health Guide for the Care and Use of Laboratory Animals (2011) and the Institutional Animal Care and Use Committees of Tulane University approved all procedures. Rats were housed individually in a temperature-controlled vivarium under a 12-h light, 12-h dark cycle and had unrestricted access to food and water. Three different cohorts of rats were used for the 3 different molecular experiments (subcellular protein fractionation, RNA extraction, and whole cell western blotting).

### 2.2. Ovariectomy and hormone treatment

Rats were anesthetized by intraperitoneal injections of ketamine (100 mg/kg ip; Bristol Laboratories, Syracuse, NY) and xylazine (7 mg/kg ip; Miles Laboratories, Shawnee, KS) and were ovariectomized. Buprenorphine (0.375 mg/kg; Reckitt Benckiser Health Care) was administered by subcutaneous injection before surgery. Ovariectomy surgery involved bilateral flank incisions through skin and muscle wall and removal of ovaries. Immediately following ovariectomy, rats were implanted with a subcutaneous 5-mm SILASTIC brand capsule (0.058 in. inner diameter and 0.077 in. outer diameter; Dow Corning, Midland, MI) on the dorsal aspect of their necks. Capsules contained either vehicle or 25% 17 $\beta$ -estradiol (Sigma-Aldrich, St. Louis, MO) diluted in vehicle. We have previously shown that implants of these dimensions and estradiol concentrations maintain blood serum estradiol

levels in middle-age retired breeders at approximately 37 pg/mL (Bohacek and Daniel, 2007), which falls within physiological range.

### 2.3. Hormone capsule replacement

Forty days after ovariectomy and capsule implantation, rats were anesthetized with ketamine and xylazine and capsules were removed and replaced with a new capsule. Buprenorphine was administered by subcutaneous injection before the start of each surgery. Rats initially receiving vehicle capsules received another vehicle capsule (Vehicle group). Rats initially receiving estradiol capsules either received a new estradiol capsule (Continuous Estradiol group) or instead received a vehicle capsule (Previous Estradiol group).

### 2.4. Euthanasia and tissue collection

Approximately 30 days following capsule replacement surgeries, rats were killed under anesthesia induced by ketamine and xylazine. Hippocampus were dissected and either quick frozen on dry ice for processing for western blotting or placed into tubes containing RNAlater (Qiagen; Hilden, Germany) for RNA extraction and stored at  $-80^{\circ}\text{C}$  until further processing.

### 2.5. Hormone treatment verification

Vaginal smears for each rat were collected for at least 4 consecutive days before capsule replacement in order to confirm hormone treatment for the initial forty-day window. Smears of ovariectomized, cholesterol-treated rats were characterized by a predominance of leukocytes, while smears of ovariectomized, estradiol-treated rats were characterized by a predominance of cornified and nucleated epithelial cells indicating hormone treatment was effective. At the time of euthanasia, a 1-cm sample of the right uterine horn was collected from each rat and weighed to verify hormone treatment during the latter part of the experiment.

### 2.6. Subcellular protein fractionation

Hippocampal tissue from each of the hormone treatment groups (Vehicle,  $n = 10$ ; Continuous Estradiol,  $n = 10$ ; Previous Estradiol,  $n = 10$ ) was lysed in cytosolic extraction buffer and protease inhibitors included in the Sub-Cellular Protein Fractionation Kit for Tissues (Thermo Scientific, Waltham, MA) using the PowerGen 125 handheld homogenizer (Fisher Scientific, San Jose, CA). Homogenate was centrifuged through the Pierce Tissue Strainer at  $500 \times g$  for 5 minutes at  $4^{\circ}\text{C}$ . Supernatant containing the cytosolic compartment extract was transferred immediately to a clean tube. The pellet was resuspended by vortexing in membrane extraction buffer containing protease inhibitors then incubated at  $4^{\circ}\text{C}$  for 10 minutes with gentle mixing. Sample was centrifuged at  $3000 \times g$  for 5 minutes at  $4^{\circ}\text{C}$ . Supernatant containing the membrane compartment extract was transferred immediately to a clean tube. The pellet was resuspended by vortexing in nuclear extraction buffer and protease inhibitors then incubated at  $4^{\circ}\text{C}$  for 30 minutes with gentle mixing. The sample was then centrifuged at  $5000 \times g$  for 5 minutes at  $4^{\circ}\text{C}$ . Supernatant containing the nuclear soluble extract was transferred to a clean tube. The pellet was resuspended by vortexing in chromatin bound extraction buffer containing room temperature nuclear extraction buffer with protease inhibitors, 100mM  $\text{CaCl}_2$ , and Micrococcal Nuclease, then incubated at  $37^{\circ}\text{C}$  for 15 minutes. The sample was then centrifuged at  $16,000 \times g$  for 5 minutes at  $4^{\circ}\text{C}$ . Supernatant containing the chromatin bound extract was added to the previously obtained nuclear soluble extract to constitute the nuclear compartment extract.

Protein concentration was determined for the cytosolic, membrane, and nuclear fractions of each sample using the Bradford Protein Assay (Thermo Scientific). Each compartment of each sample was diluted 1:1 in Laemlli Sample Buffer (BioRad, Hercules, CA) mixed with 350mM DTT (Sigma-Aldrich) and boiled for 5 minutes. One cytosolic and one membrane sample from the Vehicle group, one membrane and two nuclear samples from the Continuous Estradiol group, and one cytosolic sample from the Previous Estradiol group were excluded from western blotting either due to compartmental contamination or low protein yield.

### 2.7. Subcellular compartment western blotting

For each cytosolic, membrane, and nuclear tissue from each sample, 15 $\mu\text{g}$  of protein were loaded onto and separated on a 7.5% TGX SDS-PAGE gel at 250 V for 40 minutes. Molecular weight markers (PageRuler, Thermo Scientific) were included with each run. Proteins were transferred from gels to nitrocellulose membranes at 100 V for 30 minutes. Membranes were blocked with 5% nonfat dry milk in 1% Tween 20/1 Tris-buffered saline (TTBS) with gentle mixing at room temperature for 1 hour. After blocking, membranes were incubated with gentle mixing in primary antibody overnight at  $4^{\circ}\text{C}$  in 1% nonfat dry milk-TTBS. Samples from cytosolic, membrane, and nuclear compartments were incubated with antibodies for ER $\alpha$  (mouse monoclonal, Santa Cruz; 1:750). Samples from cytosolic fractions were incubated with antibodies for cytosolic loading control Enolase (1:2000, Santa Cruz). Samples from membrane fractions were incubated with antibodies for membrane loading control ATP1A1 (1:5000, ProteinTech). Samples from nuclear fractions were incubated with antibodies for nuclear loading control CREB (1:2000, Cell Signaling). Following primary antibody incubation, blots were washed three times for 15 minutes with TTBS. Blots were then incubated with secondary antibodies conjugated to HRP in 5% NFDM-TTBS for one hour at room temperature with gentle mixing. Secondary antibodies used were Goat Anti-Mouse IgG (1:50,000 for ER $\alpha$ , BioRad) and Goat-Anti Rabbit IgG (1:10,000 for enolase, ATP1A1, and CREB; Santa Cruz). Following secondary antibody incubation, blots were washed three times for 15 minutes with TTBS. Blots were then incubated with the chemiluminescent substrate Supersignal West Femto (Fischer Scientific) for 5 minutes and exposed to film (Kodak) for varying times to capture optimal signal intensity. Films were imaged using MCID Core imaging software (InterFocus Imaging Ltd., Cambridge, England), and optical density was measured for bands of interest. Values for each sample represent the percentage of ER $\alpha$  expression relative to the compartment-specific loading control normalized to the mean vehicle values.

### 2.8. RNA extraction

Hippocampal tissue from each of the hormone treatment groups (Vehicle,  $n=11$ ; Continuous Estradiol,  $n=11$ ; Previous Estradiol,  $n=10$ ) was homogenized using the PowerGen 125 handheld homogenizer in QIAzol Lysis Reagent (Qiagen) and extracted using the RNeasy Plus Universal Mini Kit (Qiagen). Briefly, lysate was incubated with chloroform and the aqueous phase was then incubated with ethanol. Sample was then centrifuged and washed in a spin column. RNA was eluted using RNase-free water. A gDNA eliminator was used to reduce genomic DNA contamination. RNA quality and concentration were determined by gel electrophoresis and UV absorption.

## 2.9. RT-PCR

RNA was quantified using the QuantiFast SYBR Green RT-PCR Kit (Qiagen). Primers used were for *Esr1* (QuantiTect Primer Assays; Qiagen), *Chat*, *Bdnf*, *Dlg4* and the housekeeping gene, *Gapdh* (QuantiTect Primer Assays; Qiagen). All samples were run in triplicate. Reverse transcription was performed at 50°C for 10 minutes to generate cDNA in a 25  $\mu$ L reaction volume with 50  $\mu$ g total RNA. HotStarTaq Plus DNA Polymerase was activated and reverse transcription was ended by 5 minutes of incubation at 95°C. Following the initial activation step, 40 cycles of 2-step cycling consisting of denaturation for 10 seconds at 95°C and combined annealing and/or extension for 30 seconds at 60°C were repeated. Melting curves were observed to confirm the correct primer usage in each well. Values analyzed represent a mean of triplicates normalized as a percentage of the values for the housekeeping gene, *Gapdh*.

## 2.10. Whole cell tissue processing

Hippocampal tissue from each of the hormone treatment groups (Vehicle,  $n = 8$ ; Continuous Estradiol,  $n = 8$ ; Previous Estradiol,  $n = 10$ ) were homogenized in 10  $\mu$ l/mg lysis buffer containing 1 mM EGTA, 1 mM EDTA, 20 mM Tris, 1 mM sodium pyrophosphate tetrabasic decahydrate, 4 mM 4-nitrophenyl phosphate disodium salt hexahydrate, 0.1  $\mu$ M microcystin, and 1% protease inhibitor cocktail (Sigma-Aldrich). Samples were then centrifuged for 15 minutes at 1000  $\times$  g at 4°C. Protein concentration of supernatant determined via Bradford Protein Assay. Each sample diluted 1:1 with Laemmli Sample Buffer (BioRad) mixed with 350 mM DTT, boiled for 10 minutes, and stored at -80°C until western blotting.

## 2.11. Whole cell western blotting

Twenty-five micrograms of protein were loaded onto a gel and western blotting was performed as described in Section 2.7. Primary antibodies used include PSD-95 (Millipore, 1:2000), BCL-2 (Santa Cruz, 1:2000), ChAT (Cell Signalling, 1:2000), BDNF (Santa Cruz, 1:2000), and loading control protein  $\beta$ -actin (Santa Cruz, 1:5000). Secondary antibodies conjugated to HRP were used. Blots were then incubated with the chemiluminescent substrates Supersignal West Femto (PSD-95, BDNF, ChAT) for 5 minutes or ECL Standard ( $\beta$ -actin) for 1 minute and then imaged using the ChemiDocMP Imaging System (BioRad). One sample from the Vehicle group was excluded from analysis for the BDNF western blot due to a transfer bubble over the band of interest for that sample. Optical density  $\times$  area was measured for bands of interest using MCID Imaging software. Data for each band of interest were normalized to expression of loading control protein  $\beta$ -actin.

## 2.12. Statistical analyses

Data were analyzed by One-Way ANOVA comparing treatment group and subsequent LSD post-hoc testing as appropriate. Researchers were blind to treatment group during western blotting, RT-PCR and data analysis. One extreme outlier (defined as  $\pm 3 \times$  interquartile range from the first or third quartiles for each group) was removed from the Continuous Estradiol group in the analysis of BDNF protein levels, and one extreme outlier was removed from the Previous Estradiol group in the analysis of *Dlg4* expression. All data analyses were performed using SPSS software.

## 3. Results

### 3.1. Transcriptional regulation of *Esr1* following continuous or previous estradiol exposure in the hippocampus of aging ovariectomized rats

As shown in Fig. 1A, there was no main effect of hormone treatment on hippocampal expression of *Esr1* ( $F(2,31) = 0.422$ ,  $p = 0.660$ ). These data suggest that elevated hippocampal ER $\alpha$  protein levels following continuous and previous estradiol exposure (Rodgers et al. 2010; Witty et al. 2013) during midlife are not caused by changes in transcription levels in *Esr1*.

### 3.2. Subcellular localization of ER $\alpha$ in the hippocampus of ovariectomized rats following continuous or previous exposure to estradiol in midlife

Fig. 2 displays verification of our subcellular compartment fractionation process of hippocampal tissue. Enolase—an enzyme involved in glycolysis—was used as a cytosolic marker and loading control, appearing predominately in the cytosolic fraction of samples. ATP1A1—a subunit of the sodium potassium pump ATPase—was used as a membrane marker and loading control, appearing predominately in the membrane fraction of samples. The transcription factor cAMP response element-binding protein (CREB) was used as a nuclear marker and loading control, appearing predominately in the nuclear fraction of samples. Verification of compartment fractionation was performed by western blotting for each of the compartment markers using a random selection of samples from each experimental group.


As illustrated in Fig. 1B-C, there was no main effect of hormone treatment on cytosolic ( $F(2,27) = 0.727$ ,  $p = 0.493$ ) or membrane ( $F(2,27) = 0.763$ ,  $p = 0.477$ ) ER $\alpha$  levels. As illustrated in Fig. 1D, there was a main effect of hormone treatment on nuclear ER $\alpha$  levels ( $F(2,27) = 3.396$ ,  $p = 0.0496$ ), with levels increased in both the Continuous Estradiol ( $p = 0.033$ ) and Previous Estradiol ( $p = 0.036$ ) groups as compared to Vehicle group levels. There was no significant difference in nuclear ER $\alpha$  levels between the Continuous Estradiol and Previous Estradiol groups ( $p = 0.863$ ). These results demonstrate that previous exposure to estradiol in midlife results in lasting elevation specifically of nuclear ER $\alpha$  levels in the hippocampus of ovariectomized rats, similar to levels in animals receiving ongoing estradiol treatment.

### 3.3. Hippocampal transcriptional regulation and protein expression of genes that contain ERE sequences following continuous or previous midlife estradiol exposure

As illustrated in Fig. 3A, there was a main effect of hormone treatment on expression of *Bdnf* ( $F(2,31) = 4.180$ ,  $p = 0.025$ ), with increased RNA levels in both the Continuous Estradiol ( $p = 0.011$ ) and Previous Estradiol ( $p = 0.039$ ) groups as compared to the levels in the Vehicle group. No significant difference in RNA levels of *Bdnf* was found between the Continuous Estradiol and Previous Estradiol groups ( $p = 0.621$ ). There was a main effect of hormone treatment on hippocampal protein levels of BDNF (Fig. 3B,  $F(2,23) = 8.580$ ,  $p = 0.002$ ), with levels in the Previous Estradiol group significantly increased as compared to those in both the Vehicle ( $p = 0.018$ ) and the Continuous Estradiol ( $p = 0.001$ ). No significant difference in protein levels of BDNF was found between the Vehicle and Continuous Estradiol groups ( $p = 0.195$ ).

As illustrated in Fig. 3C, there was a main effect of hormone treatment on expression of *Chat* (Fig. 2C,  $F(2,31) = 4.810$ ,





**Fig. 1.** Transcriptional Regulation of *Esr1* and subcellular localization of ER $\alpha$  in the hippocampus of ovariectomized rats following continuous or previous exposure to estradiol in midlife. Middle-aged female rats were ovariectomized and treated to one of 3 hormone conditions via Silastic capsule: Vehicle (Veh), which received a vehicle capsule; Continuous Estradiol (Cont E), which received an estradiol capsule for the duration of the experiment; or Previous Estradiol (Prev E), which received an estradiol capsule for 40 days followed by a vehicle capsule for the duration of the experiment. One month later, hippocampi were processed for either RNA extraction and RT-PCR using primers for *Esr1* and housekeeping gene *Gapdh* or for subcellular fractionation and western blotting for ER $\alpha$ , measured by density x area of ER $\alpha$ /loading control proteins. (A) There was no significant effect of hormone treatment on *Esr1* expression in the hippocampus relative to *Gapdh* expression. (B-C) There was no significant effect of hormone treatment on cytosolic (B) or membrane (C) ER $\alpha$ . (D) There was a main effect of hormone treatment on levels of nuclear ER $\alpha$  ( $p < 0.05$ ). Post hoc testing revealed increased levels in the Cont E and Prev E groups relative to the Veh group. Data are presented as means  $\pm$  SEM normalized to percent Vehicle group. \* $p < 0.05$  versus Veh.

$p = 0.016$ ), with increased RNA levels in both the Continuous Estradiol ( $p = 0.006$ ) and Previous Estradiol ( $p = 0.035$ ) groups as compared to the levels in the Vehicle group. No significant difference in RNA levels of *Chat* was found between the Continuous Estradiol and Previous Estradiol groups ( $p = 0.492$ ). There was a main effect of hormone treatment on hippocampal protein levels of ChAT (Fig. 3D,  $F(2,25) = 5.944$ ,  $p = 0.008$ ), with levels in the Continuous Estradiol ( $p = 0.005$ ) and Previous Estradiol ( $p = 0.007$ ) groups significantly increased as compared to those in the Vehicle group. No significant difference in ChAT protein levels of was found between the Continuous Estradiol and Previous Estradiol groups ( $p = 0.770$ ).


#### 3.4. Hippocampal transcriptional regulation and protein expression of one gene that does not contain an ERE sequence following continuous or previous midlife estradiol exposure

As illustrated in Fig. 4A, there was a main effect of hormone treatment on expression of *Dlg4* ( $F(2,30) = 4.726$ ,  $p = 0.017$ ), with increased RNA levels in both the Continuous Estradiol ( $p = 0.006$ ) and Previous Estradiol ( $p = 0.043$ ) groups as compared to the

levels in the Vehicle group. No significant difference in RNA levels of *Dlg4* was found between the Continuous Estradiol and Previous Estradiol groups ( $p = 0.496$ ). There was no main effect of hormone treatment on hippocampal protein levels of PSD-95 ( $F(2,25) = 0.088$ ,  $p = 0.916$ ).

#### 4. Discussion

Results of the present studies reveal that previous exposure to estradiol during midlife has lasting impacts on hippocampal function through sustained transcriptional activity of ER $\alpha$  that persists long after estradiol treatment has ended. Previous estradiol exposure resulted in lasting increases in the nuclear pool of hippocampal ER $\alpha$  as well as long-term upregulation of BDNF and ChAT in aging ovariectomized rats—effects that were similar to those observed in ovariectomized animals with ongoing estradiol exposure. Specifically, we showed that both continuous and previous estradiol increased nuclear ER $\alpha$  protein levels in the hippocampus of aging, ovariectomized rats one month after cessation of estradiol treatment. Animals from the Previous Estradiol group also had elevated hippocampal expression of 2 ERE-dependent genes



**Fig. 2.** Verification of subcellular compartment fractionation. Hippocampal tissue was processed for subcellular fractionation in order to separate the cytosolic, membrane, and nuclear compartments of cells via consecutive centrifugation steps using a commercially available kit. Compartment separation was verified using western blotting for cytosolic marker enolase, membrane marker ATP1A1, and nuclear marker CREB on samples from all compartments.

(*Bdnf*, *Chat*) and their corresponding proteins (BDNF, ChAT), as well as elevated expression of one non-ERE-dependent gene (*Dlg4*), as compared to animals from the Vehicle group. Animals receiving ongoing estradiol exposure displayed similar, but not identical, expression patterns in the hippocampus, with elevated gene expression of *Bdnf*, *Chat*, and *Dlg4*, but only elevated protein levels of ChAT as compared to animals from the Vehicle group. Overall, these findings indicate a crucial role for ER $\alpha$  in maintaining hippocampal memory in ovariectomized animals through sustained transcriptional activity of the receptor following previous estradiol exposure in midlife in a pattern that is comparable to that observed in the presence of circulating estradiol.


Maintained hippocampal protein levels of ER $\alpha$  following continuous and previous estradiol exposure in ovariectomized rats has been associated with enhanced hippocampal-dependent memory (Rodgers et al., 2010). As a nuclear steroid hormone receptor, ER $\alpha$  can act in a variety of functions within a cell. Here we demonstrated through subcellular compartment fractionation that ER $\alpha$  is specifically increased in the nuclear compartment of hippocampal cells following continuous or previous midlife estradiol exposure in ovariectomized rats, although the receptor is also present in the cytosol and membrane compartments in all hormone treatment groups. This finding aligns with recent work from our lab demonstrating that short-term estradiol treatment immediately after ovariectomy results in lasting enhancements in hippocampal memory and increased ERE-dependent transcriptional activity in mice (Pollard et al., 2018). Together, it strengthens the connection between maintained protein levels of ER $\alpha$  following previous exposure to estradiol in midlife and enhanced memory via maintained transcriptional activity of ER $\alpha$  in the hippocampus.

In contrast to the increase induced by continuous and previous estradiol treatments on protein levels of nuclear ER $\alpha$  in the hippocampus, we saw no effect of either estradiol treatment on *Esr1* transcription levels. These results were somewhat unexpected because of the presence of an ERE on the gene *Esr1*. There remains much disagreement in the literature as to the impact of ovariectomy and estradiol treatment on *Esr1* mRNA levels throughout the brain, with changes in hippocampal *Esr1* mRNA levels vary-

ing based on age, length of time following ovariectomy, and sub-region of the hippocampus examined (Ivanov et al., 2017). However, besides modification of ER $\alpha$  gene transcription, ER $\alpha$  levels can be impacted by changes in receptor degradation rate. ER $\alpha$  is degraded via the ubiquitin-proteasomal pathway (Tateishi et al., 2004). When the receptor is unliganded, the E3 ubiquitin ligase, C terminus of Hsc70-interacting protein (CHIP) binds ER $\alpha$  and targets it for ubiquitination and proteasomal degradation (Fan et al., 2005). Long-term estrogen deprivation following OVX increases interaction between CHIP and ER $\alpha$  and thus subsequent ubiquitination of ER $\alpha$  (Zhang et al., 2011). Prior results from our lab indicate that previous estradiol treatment may prevent effects of estrogen deprivation on ER $\alpha$  degradation. The same 40 days of previous estradiol treatment used in the current study resulted in lasting decreased association between ER $\alpha$  and CHIP in parallel to increased protein levels of ER $\alpha$  when measured 1 month following termination of the previous estradiol treatment (Black et al., 2016). Together, these results with those of the current work provide support for the hypothesis that lasting changes in levels of ER $\alpha$  resulting from previous exposure to midlife estradiol are primarily due to lasting changes in protein degradation rather than transcription.

The ability of estradiol treatment to increase levels of ER $\alpha$  specifically in the nucleus is consistent with its ability to impact estradiol-sensitive genes and proteins. We found impacts of both continuous and previous estradiol treatments on hippocampal gene expression and protein levels in 2 ERE-dependent genes known to impact memory. Animals that had previously been exposed to estradiol following ovariectomy displayed upregulated expression two *Bdnf* and *Chat* in the hippocampus one month after termination of estradiol exposure. These increases in gene expression observed in the Previous Estradiol group were comparable to those found in the hippocampi of animals receiving ongoing estradiol exposure, demonstrating that short-term estradiol exposure during midlife can have lasting effects on hippocampal gene expression. There were, however, differences between the Continuous Estradiol and Previous Estradiol group in the expression patterns of the proteins associated with these genes.

Significant increases in hippocampal protein levels of both BDNF and ChAT were found in the Previous Estradiol group, consistent with the observed increases in gene expression of *Bdnf* and *Chat* described above. These findings were also consistent with earlier findings from our lab that previous estradiol exposure during midlife results in lasting increases in levels of ChAT in the hippocampus (Rodgers et al., 2010; Witty et al., 2013) but which has not been shown before for hippocampal BDNF levels. In contrast to the effects observed in the Previous Estradiol group, animals in the Continuous Estradiol group showed significantly increased hippocampal protein levels of ChAT, but not BDNF, despite showing increased gene expression for both *Bdnf* and *Chat*. Several earlier studies have demonstrated that estradiol exposure increases *Chat* mRNA (Gibbs, 1996), elevates ChAT protein expression (Gibbs, 1997; Rodgers et al., 2010) and increases acetylcholine release in the hippocampus of ovariectomized rodents (Gabor et al., 2003; Gibbs, 1997). However, the relationship between estradiol exposure, *Bdnf* mRNA, and BDNF protein levels in ovariectomized animals is less clear. Consistent with our findings, several—but not all (Cavus and Duman, 2003)—studies found continuous estradiol treatment to increase hippocampal mRNA levels of *Bdnf* in ovariectomized animals (Berchtold et al., 2001; Liu et al., 2001; Singh et al., 2005). There is however no clear consensus as to the effect of estradiol treatment on hippocampal BDNF protein levels, with some studies showing increased protein expression throughout the hippocampus (Berchtold et al., 2001), some showing increased BDNF levels only in specific subregions of the hippocampus (Zhou et al., 2005), and others showing decreased hippocampal protein levels in ovariectomized animals treated with estra-




**Fig. 3.** Hippocampal transcriptional regulation and protein expression of genes that contain ERE sequences following continuous or previous midlife estradiol exposure. Middle-aged female rats were ovariectomized and treated to one of 3 hormone conditions via Silastic capsule: Vehicle (Veh), which received a vehicle capsule; Continuous Estradiol (Cont E), which received an estradiol capsule for the duration of the experiment; or Previous Estradiol (Prev E), which received an estradiol capsule for 40 days followed by a vehicle capsule for the duration of the experiment. One month later, hippocampi were processed for RNA extraction and RT-PCR using primers for *Bdnf*, *Chat*, and *Gapdh*, or for western blotting for BDNF, ChAT, and  $\beta$ -actin. RT-PCR data were normalized to housekeeping gene *Gapdh* and western blot data were normalized to loading control protein  $\beta$ -actin. (A–B) There was a main effect of hormone treatment ( $p < 0.05$ ) on *Bdnf* RNA expression (A) and BDNF protein levels (B) in the hippocampus. Post hoc testing revealed increased *Bdnf* expression in the Cont E and Prev E groups and increased BDNF protein levels in the Prev E group as compared to the Veh group. (C–D) There was a main effect of hormone treatment ( $p < 0.05$ ) on *Chat* RNA expression (C) and ChAT protein levels (D) in the hippocampus. Post hoc testing revealed increased *Chat* expression and increased ChAT protein levels in the Cont E and Prev E groups as compared to the Veh group. \* $p < 0.05$  versus Veh.

diol (Gibbs, 1999). The results of the current studies therefore add to this complex story by demonstrating that previous exposure to estradiol during midlife results in lasting increases in hippocampal mRNA and protein expression of BDNF, whereas continuous exposure to estradiol only results in increased mRNA levels. Nevertheless, both hormone treatments result in lasting increases in mRNA and protein levels of ChAT, another ERE-dependent gene critical for memory. As described above regarding estrogenic regulation of *Esr1* mRNA, the relationship between estrogen receptor activity, mRNA expression, and protein levels can vary greatly due to brain region, age, and duration of estrogen exposure, among other variables (Scharfman and MacLuskey, 2005).

Finally, our findings demonstrate that continuous or previous estradiol exposure following ovariectomy can also result in increased transcription of non-ERE-dependent genes, with both the Continuous Estradiol and Previous Estradiol groups showing increased hippocampal expression of *Dlg4* as compared to the Vehicle group. Interestingly, there was no observed change in levels of PSD-95, the protein transcribed by *Dlg4*, in either hor-

mon treatment group. Estradiol treatment has repeatedly been shown to increase protein levels of PSD-95 in the hippocampus (Nelson et al., 2014; Waters et al., 2009), a rapid effect that is attributed to the PI3K-Akt signaling pathway activated by membrane ER $\alpha$  (Akama and McEwen, 2003; Murakami et al., 2015). Because we observed no change in membrane levels of ER $\alpha$  following continuous or previous estradiol treatments, it is unclear how increased ER $\alpha$  in the nucleus of hippocampal cells can result in lasting changes in gene expression but not protein levels of PSD-95. Future studies should investigate potential crosstalk between membrane and nuclear ER $\alpha$  and its impact on synaptic proteins following previous midlife estradiol exposure.

Together, the results of the present study indicate a critical role for ER $\alpha$  as a transcriptional regulator of hippocampal function in both the presence and absence of circulating estrogens. Previous exposure to estradiol exposure in midlife results in lasting increases in nuclear ER $\alpha$  activity in the hippocampus, resulting in increased transcription of genes important for hippocampal function and enhanced hippocampal dependent memory (Rodgers et al.,



**Fig. 4.** Hippocampal transcriptional regulation and protein expression of gene that does not contain an ERE sequence following continuous or previous midlife estradiol exposure. Middle-aged female rats were ovariectomized and treated to one of 3 hormone conditions via Silastic capsule: Vehicle (Veh), which received a vehicle capsule; Continuous Estradiol (Cont E), which received an estradiol capsule for the duration of the experiment; or Previous Estradiol (Prev E), which received an estradiol capsule for 40 days followed by a vehicle capsule for the duration of the experiment. One month later, hippocampi were processed for RNA extraction and RT-PCR using primers for *Dlg4* and *Gapdh*, or for western blotting for PSD-95 and  $\beta$ -actin. RT-PCR data were normalized to housekeeping gene *Gapdh* and western blot data were normalized to loading control protein  $\beta$ -actin. (A) There was a main effect of hormone treatment ( $p < 0.05$ ) on *Dlg4* RNA expression in the hippocampus. Post hoc testing revealed increased *Dlg4* expression in the Cont E and Prev E groups as compared to the Veh group. (B) There was no effect of hormone treatment on PSD-95 protein levels. \* $p < 0.05$  versus Veh.

2010). In the current work, we assessed effects of previous estradiol treatment on our measures one month following termination of the treatment. However, we expect that these effects persist beyond the 1-month period as we have previously demonstrated that this same model of 40 days of midlife estradiol exposure results in similar enhancements in memory and increased protein levels of ER $\alpha$  at 1, 2.5, and 7 months after termination of treatment in rats (Black et al., 2016; Rodgers et al., 2010; Witty et al., 2013), the later time frame equivalent to decades in humans. Thus, the current findings demonstrate a mechanism through which long-term changes in memory can occur by sustained activation of nuclear ER $\alpha$  in the hippocampus. Future work should thoroughly examine the mechanisms through which ER $\alpha$  can influence hippocampal function in the absence of circulating estrogens, though previous work from our lab indicates a role for brain-derived neuroestrogens (Baumgartner et al., 2019) as well as ligand-independent activation of ER $\alpha$  by growth factors including insulin-like growth factor-1 (IGF-1) (Grissom and Daniel, 2016; Witty et al., 2013).

Ultimately these findings have important implications for women who use short-term estrogen treatment to treat their menopause symptoms. Several studies, including those of women who underwent surgical menopause earlier in life than natural menopause (Bove et al., 2014; Rocca et al., 2014), suggest a lasting benefit for cognition of short-term estrogen use immediately following menopause (Bagger et al., 2005; Whitmer et al., 2011). Future work should expand upon the clinical relevance of this model of short-term estrogen use for women undergoing natural menopause, in which the loss of ovarian hormones is less abrupt and estrogen treatment is often paired with progestins that may have different impacts on estrogen receptor function in the brain. Ongoing large-scale clinical studies such as the Kronos Early Estrogen Prevention Study (KEEPS) will provide more insight into the long-term impacts of short-term estrogen use in midlife for women experiencing natural menopause (Wharton et al., 2013). Nevertheless, the results of the current study highlight a potential mechanism for short-term midlife estrogen use to have lasting impacts on cognition by maintaining transcriptional activity of nuclear ER $\alpha$  in the hippocampus. Furthermore, these findings emphasize the critical impact that ER $\alpha$  can have on the aging female brain in the absence of circulating estrogens.

#### Submission declaration and verification

I certify that this manuscript is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder.

#### Disclosure statement

None.

#### Acknowledgements

This work was supported by the National Institute on Aging under Grant [RF1AG041374](#)

#### Authors' contributions

Conceptualization and experiment planning by NEB, KLB, and JMD. Data collection and analysis by NEB and SMM. Writing by NEB and JMD.

#### Credit author statement

**Nina Baumgartner:** Conceptualization, Methodology, Investigation, Analysis, Writing. **Katelyn Black:** Conceptualization, Methodology, Validation. **Shannon McQuillen:** Investigation, Analysis. **Jill M. Daniel:** Conceptualization, Methodology, Writing, Supervision, Funding acquisition.

#### References

- Akama, K., McEwen, B., 2003. Estrogen stimulates postsynaptic density-95 rapid protein synthesis via the Akt/protein kinase B pathway. *J Neurosci* 23, 2333–2339.
- Bagger, Y., Tankó, L., Alexandersen, P., Qin, G., Christiansen, C., 2005. Early postmenopausal hormone therapy may prevent cognitive impairment later in life. *Menopause* 12, 12–17.



- Baumgartner, N., Grissom, E., Pollard, K., McQuillen, S., Daniel, J., 2019. Neuroestrogen-dependent transcriptional activity in the brains of ERE-Luciferase reporter mice following short- and long-term ovariectomy. *eNeuro* 6, 1–11 ENEURO.0275–0219.2019.
- Baxter, M., Santistevan, A., Bliss-Moreau, E., Morrison, J., 2018. Timing of cyclic estradiol treatment differentially affects cognition in aged female rhesus monkeys. *Behav Neurosci* 132, 213–223.
- Berchtold, N., Kesslak, J., Pike, C., Adlard, P., Cotman, C., 2001. Estrogen and exercise interact to regulate brain-derived neurotrophic factor mRNA and protein expression in the hippocampus. *Eur J Neurosci* 14, 1992–2002.
- Bi, R., Foy, M., Thompson, R., Baudry, M., 2003. Effects of estrogen, age, and calpain on MAP kinase and NMDA receptors in female rat brain. *Neurobiol Aging* 24, 977–983.
- Black, K.L., Baumgartner, N.E., Daniel, J.M., 2018. Lasting impact on memory of midlife exposure to exogenous and endogenous estrogens. *Behav. Neurosci.* 132, 547–551.
- Black, K.L., Witty, C.F., Daniel, J.M., 2016. Previous midlife oestradiol treatment results in long-term maintenance of hippocampal oestrogen receptor alpha levels in ovariectomized rats: mechanisms and implications for memory. *J Neuroendocrinol* 28, 1–10.
- Bohacek, J., Daniel, J.M., 2007. Increased daily handling of ovariectomized rats enhances performance on a radial-maze task and obscures effects of estradiol replacement. *Horm Behav* 52, 237–243.
- Bove, R., Secor, E., Chibnik, L., Barnes, L., Schneider, J., Bennett, D., De Jager, P., 2014. Age at surgical menopause influences cognitive decline and Alzheimer pathology in older women. *Neurology* 82, 222–229.
- Castles, C., Oesterreich, S., Hansen, R., Fugua, S., 1997. Auto-regulation of the estrogen receptor promoter. *Ster Biochem Mol Bio* 62, 155–163.
- Cavus, I., Duman, R., 2003. Influence of estradiol, stress, and 5-HT<sub>2A</sub> agonist treatment on brain-derived neurotrophic factor expression in female rats. *Biol. Psychiatry* 54, 59–69.
- Chakraborty, T., Gore, A., 2004. Aging-related changes in ovarian hormones, their receptors, and neuroendocrine function. *Exp Biol Med* 229, 977–987.
- Chen, W., Manson, J., Hankinson, S., Rosner, B., Holmes, M., Willett, W., Colditz, G., 2006. Unopposed estrogen therapy and the risk of invasive breast cancer. *Arch Intern Med* 166, 1027–1032.
- Daniel, J.M., Hulst, J.L., Berbling, J.L., 2006. Estradiol replacement enhances working memory in middle-aged rats when initiated immediately after ovariectomy but not after a long-term period of ovarian hormone deprivation. *Endocrinology* 147, 607–614.
- Daniel, J.M., Witty, C.F., Rodgers, S.P., 2015. Long-term consequences of estrogens administered in midlife on female cognitive aging. *Horm Behav* 74, 77–85.
- Fan, M., Park, A., Nephew, K., 2005. CHIP (carboxyl terminus of Hsc70-interacting protein) promotes basal and geldanamycin-induced degradation of estrogen receptor- $\alpha$ . *Mol Endocrinol* 19, 2901–2914.
- Foster, T., 2012. Role of estrogen receptor alpha and beta expression and signaling on cognitive function during aging. *Hippocampus* 22, 656–669.
- Foy, M., Baudry, M., Foy, J., Thompson, R., 2008. 17 $\beta$ -estradiol modifies stress-induced and age-related changes in hippocampal synaptic plasticity. *Behav Neurosci* 122, 301–309.
- Fugger, H., Kumar, A., Lubahn, D., Korach, K., Foster, T., 2001. Examination of estradiol effects on the rapid estradiol mediated increase in hippocampal synaptic transmission in estrogen receptor alpha knockout mice. *Neurosci Lett* 309, 207–209.
- Gabor, R., Nagle, R., Johnson, D., Gibbs, R., 2003. Estrogen enhances potassium-stimulated acetylcholine release in the rat hippocampus. *Brain Res* 962, 244–247.
- Gibbs, R., 1996. Fluctuations in relative levels of choline acetyltransferase mRNA in different regions of the rat basal forebrain across the estrous cycle: effects of estrogen and progesterone. *J Neuro* 16, 1049–1055.
- Gibbs, R., 1997. Effects of estrogen on basal forebrain cholinergic neurons vary as a function of dose and duration of treatment. *Brain Res* 757, 10–16.
- Gibbs, R., 1999. Treatment with estrogen and progesterone affects relative levels of brain-derived neurotrophic factor mRNA and protein in different regions of the adult rat brain. *Brain Res* 844, 20–27.
- Grissom, E.M., Daniel, J.M., 2016. Evidence for ligand-independent activation of hippocampal estrogen receptor- $\alpha$  by IGF-1 in hippocampus of ovariectomized rats. *Endocrinology* 157, 3149–3156.
- Hyder, S., Chiappetta, C., Stancel, G., 1999. Interaction of human estrogen receptors alpha and beta with the same naturally occurring estrogen response elements. *Biochem Pharmacol* 57, 597–601.
- Ianov, L., De Both, M., Chawla, M.K., Rani, A., Kennedy, A.J., Piras, I., Day, J.J., Siniard, A., Kumar, A., Sweatt, J.D., Barnes, C.A., Huentelman, M.J., Foster, T.C., 2017. Hippocampal Transcriptomic Profiles: Subfield Vulnerability to Age and Cognitive Impairment. *Front Aging Neurosci* 9 (383), 1–15.
- Klinge, C., 2001. Estrogen receptor interaction with estrogen response elements. *Nucleic Acids Res* 29, 2905–2919.
- Koebele, S.V., Bimonte-Nelson, H.A., 2017. The endocrine-brain-aging triad where many paths meet: female reproductive hormone changes at midlife and their influence on circuits important for learning and memory. *Exp Gerontol* 94, 14–23.
- Közszegi, Z., Szego, É., Cheong, R., Tolod-Kemp, E., Ábrahám, I., 2011. Postlesion estradiol treatment increases cortical cholinergic innervations via estrogen receptor- $\alpha$  dependent nonclassical estrogen signaling *in Vivo*. *Endocrinology* 152, 3471–3482.
- Kumar, A., Foster, T., 2002. 17 $\beta$ -estradiol benzoate decreases the AHP amplitude in CA1 pyramidal neurons. *J Neurophysiol* 88, 621–626.
- Liu, Y., Fowler, C., Young, L., Yan, Q., Insel, T., Wang, Z., 2001. Expression and estrogen regulation of brain-derived neurotrophic factor gene and protein in the forebrain of female prairie voles. *J Comp Neuro* 433, 499–514.
- Luine, V., 1985. Estradiol increases choline acetyltransferase activity in specific basal forebrain nuclei and projection areas of female rats. *Exp Neurol* 89, 484–490.
- Luine, V., Frankfurt, M., 2020. Estrogenic regulation of memory: the first 50 years. *Hormones and Behavior* 121, 104711.
- Ma, S., Tang, N., Leung, G., Fung, A., Lam, L., 2014. Estrogen receptor  $\alpha$  polymorphisms and the risk of cognitive decline: a 2-year follow-up study. *Am J Geriatr Psychiatry* 22, 489–498.
- Ma, S., Tang, N., Tam, C., Lui, V., Lau, E., Zhang, Y., Chiu, H., Lam, L., 2009. Polymorphisms of the estrogen receptor alpha (ESR1) gene and the risk of Alzheimer's disease in a southern Chinese community. *Int Psychogeriatr* 21, 977–986.
- Maki, P., Dennerstein, L., Clark, M., Guthrie, J., LaMontagne, P., Fornelli, D., Little, D., Henderson, V., Resnick, S., 2011. Perimenopausal use of hormone therapy is associated with enhanced memory and hippocampal function later in life. *Brain Res* 1379, 232–243.
- Murakami, G., Hojo, Y., Ogiue-Ikeda, M., Mukai, H., Chambon, P., Nakajima, K., Ooishi, Y., Kimoto, T., Kawato, S., 2015. Estrogen receptor KO mice study on rapid modulation of spines and long-term depression in the hippocampus. *Brain Res* 1621, 133–146.
- Nelson, B., Springer, R., Daniel, J., 2014. Antagonism of brain insulin-like growth factor-1 receptors blocks estradiol effects on memory and levels of hippocampal synaptic proteins in ovariectomized rats. *Psychopharmacology (Berl)* 231, 899–907.
- Pencea, V., Bingaman, K., Wiegand, S., Luskin, M., 2001. Infusion of brain-derived neurotrophic factor into the lateral ventricle of the adult rat leads to new neurons in the parenchyma of the striatum, septum, thalamus, and hypothalamus. *J Neurosci* 21, 6706–6717.
- Pollard, K.J., Wartman, H.D., Daniel, J.M., 2018. Previous estradiol treatment in ovariectomized mice provides lasting enhancement of memory and brain estrogen receptor activity. *Horm Behav* 102, 76–84.
- Rocca, W., Grossardt, B., Shuster, L., 2014. Oophorectomy, estrogen, and dementia: a 2014 update. *Mol Cell Endocrinol* 389, 7–12.
- Rodgers, S.P., Bohacek, J., Daniel, J.M., 2010. Transient estradiol exposure during middle age in ovariectomized rats exerts lasting effects on cognitive function and the hippocampus. *Endocrinology* 151, 1194–1203.
- Ryan, J., Carrière, I., Carcaillon, L., Dartigues, J., Auriaucombe, S., Rouaud, O., Berr, C., Ritchie, K., Scarabin, P., Ancelin, M., 2014. Estrogen receptor polymorphisms and incident dementia: the prospective 3C study. *Alzheimers Dement* 10, 27–35.
- Santen, R., Allred, D., Ardoin, S., Archer, D., Boyd, N., Braunstein, G., Burger, H., Colditz, G., Davis, S., Gambacciani, M., Gower, B., Henderson, V., Jarjour, W., Karas, R., Kleerekoper, M., Lobo, R., Manson, J., Marsden, J., Martin, K., Martin, L., Pinkerton, J., Rubinow, D., Teede, H., Thiboutot, D., Utian, W., 2010. Postmenopausal hormone therapy: an endocrine society scientific statement. *Clin Endocrinol Metab* 95, s1–s66.
- Scharfman, H., MacLuskey, N., 2005. Similarities between actions of estrogen and BDNF in the hippocampus: coincidence or clue? *Trends in Neurosci* 28, 79–85.
- Singh, M., Meyer, E., Simpkins, J., 2005. The effect of ovariectomy and estradiol replacement on brain-derived neurotrophic factor messenger ribonucleic acid expression in cortical and hippocampal brain regions of female Sprague-Dawley rats. *Endocrinology* 136, 2320–2324.
- Sohrabji, F., Lewis, D., 2006. Estrogen-BDNF interactions: implications for neurodegenerative diseases. *Front Neuroendocrinol* 27, 404–414.
- Sohrabji, F., Miranda, R., Toran-Allerand, C., 1995. Identification of a putative estrogen response element in the gene encoding brain-derived neurotrophic factor. *Proc Natl Acad Sci USA* 92, 11110–11114.
- Tateishi, Y., Kawabe, Y., Chiba, T., Murata, S., Ichikawa, K., Murayama, A., Tanaka, K., Baba, T., Kato, S., Yanagisawa, J., 2004. Ligand-dependent switching of ubiquitin-proteasome pathways for estrogen receptor. *EMBO J* 23, 4813–4823.
- Waters, E., Mitterling, K., JL, S., Mazid, S., McEwen, B., Milner, T., 2009. Estrogen receptor alpha and beta specific agonists regulate expression of synaptic proteins in rat hippocampus. *Brain Res* 1290, 1–11.
- Wharton, W., Gleason, C., Miller, V., Asthana, S., 2013. Rationale and design of the Kronos Early Estrogen Prevention Study (KEEPS) and the KEEPS Cognitive and Affective sub study (KEEPS Cog). *Brain Res* 1514, 12–17.
- Whitmer, R., Quesenberry, C., Zhou, J., Yaffe, K., 2011. Timing of hormone therapy and dementia: the critical window theory revisited. *Ann Neurol* 69, 163–169.
- Witty, C., Foster, T., Semple-Rowland, S., Daniel, J., 2012. Increasing hippocampal estrogen receptor alpha levels via viral vectors increases MAP kinase activation and enhances memory in aging rats in the absence of ovarian estrogens. *PLoS One* 7, e51385.
- Witty, C., Gardella, L., Perez, M., Daniel, J., 2013. Short-term estradiol administration in aging ovariectomized rats provides lasting benefits for memory and the hippocampus: a role for insulin-like growth factor-1. *Endocrinology* 154, 842–852.
- Yaffe, K., Lindquist, K., Sen, S., Cauley, J., Ferrell, R., Penninx, B., Harris, T., Li, R., Cummings, S., 2009. Estrogen receptor genotype and risk of cognitive impairment in elders: findings from the Health ABC study. *Neurobiol Aging* 30, 607–614.
- Zhang, Q.G., Han, D., Wang, R.M., Dong, Y., Yang, F., Vadlamudi, R.K., Brann, D.W., 2011. C terminus of Hsc70-interacting protein (CHIP)-mediated degradation of hippocampal estrogen receptor- $\alpha$  and the critical period hypothesis of estrogen neuroprotection. *Proc Natl Acad Sci USA* 108, E617–E624.
- Zhou, J., Zhang, H., Cohen, R., Pandey, S., 2005. Effects of estrogen treatment on expression of brain-derived neurotrophic factor and cAMP response element-binding protein expression and phosphorylation in rat amygdaloid and hippocampal structures. *Neuroendocrinology* 81, 294–310.